

JETTER

Bispyribac Sodium 10% W/V SC

JETTER controls major grasses, sedges and broad leaf weeds of rice. It has wide application window from 2-5 leaf stages of weeds.

JETTER Suitable for need based application only when the weeds emerge.

JETTER is safe to rice due to its selectivity.

JETTER gets absorbed quickly in weeds, rain fastness is 6 hours of application.

JETTER has a low dosage of 80-120 ml/acre.

Method of Application:

Water in the field should be drained out at the time of spray and has to re-flood the field within 48 - 72 hrs. of application. There after maintain water in the field for 5-7 days to suppress weed emergence. Shake the bottle well before use. Targeted weeds should directly be exposed to **JETTER** by ensuring uniform spray. Use Flat fan / Flood jet nozzle for spraying the Herbicides



CROP	WEEDS	Dosage MI/Ha	Dilution in Water (Litres/Ha)
Rice (Nursery)	Echinochloa crusgalli, Echinochloa colonum, Ischaemum rugosum, Cyperus difformis,	200	300
Rice (Transplanted)	Cyperus iria, Fimbristylis miliacea, Eclipta alba, Ludwigia parviflora,	200-250	300
Rice (Directly Seeded)	Monochoria vaginalis, Alternanthera philoxeroides, Sphenoclea zeylenica.	200-250	300

Pack size : 10ml, 50ml, 100ml, 200 ml, 500ml