

MAYOO-C

Carbendazim 12%+ Mancozeb 63% WP

MAYOO-C Broad Spectrum Systemic and contact Fungicide

MAYOO-C is effective prophylactic & curative fungicide.

MAYOO-C ensures double protection from systemic & contact action.

MAYOO-C controls leaf spot & rust diseases of groundnut and blast disease of Rice crop.

MAYOO-C effectively controls fungal diseases by systemic and contact action.

MAYOO-C is used as seed treatment, soil drenching, rhizome / tuber dip and foliar sprays.

MAYOO-C acts by inhibiting the development of fungal germ tube, formation of appressoria and growth of mycelia, also cause disturbance in fungal enzyme functioning.

MAYOO-C is effective against a wide range of fungal diseases in various crops.



Crop	Disease	Dosage / Ha		Waiting period
		Formulation (Gm)	Dilution in water (litres)	
Groundnut	Leaf spot, Rust	500	500	72 days
Rice	Blast	750	750	57 days
Chilli	Leaf spot, Fruit rot, Powdery mildew	750	500	3 days
Grape	Downy mildew, Powdery mildew, Anthracnose	0.15%	As required depends upon crop canopy	7 days
Mango	Powdery mildew, Anthracnose	0.15%	As required depends upon crop canopy	7 days
Potato	Early blight, Late blight, Black scurf	1750	500	47 days
Tea	Blister blight, Gray blight, Red rust, Die back, Black rot	1250	250-500	7 days

PACK SIZE : 100g, 250g, 500g & 1Kg.